

Metadynamics Remedies for Topological Freezing

Francesco Sanfilippo

UNIVERSITY OF
Southampton
School of Physics
and Astronomy

(that's where you'll come at the end of July)

Mainly based on arXiv:1508.07270

“Metadynamics Surfing on Topology Barriers: the $CP(N - 1)$ Case”

A.Laio, G.Martinelli, F.S



The Illness

- 1 Topological charge
- 2 Critical Slowing Down



The Illness

- 1 Topological charge
- 2 Critical Slowing Down



The Treatment

- 1 Metadynamics
- 2 A case of investigation: $CP(N-1)$ model



The Illness

- 1 Topological charge
- 2 Critical Slowing Down



The Treatment

- 1 Metadynamics
- 2 A case of investigation: $CP(N-1)$ model



Side Effects (and side outcomes!)

- 1 Measuring the Free Energy
- 2 Reweighting



The Illness

- 1 Topological charge
- 2 Critical Slowing Down



The Treatment

- 1 Metadynamics
- 2 A case of investigation: $CP(N-1)$ model



Side Effects (and side outcomes!)

- 1 Measuring the Free Energy
- 2 Reweighting



Extension and perspectives

- 1 First checks in QCD
- 2 Extension of the method

Homotopy group

Topological sector: set of configurations that can be transformed one into the other by means of a continuous deformation

Topological charge

Homotopy group

Topological sector: set of configurations that can be transformed one into the other by means of a continuous deformation

Winding number

Topological charge **density** in QCD

$$q(x) = \frac{1}{32\pi^2} \epsilon_{\mu\nu\rho\sigma} \text{Tr} [F_{\mu\nu}(x) F_{\rho\sigma}(x)]$$

- Its volume integral defines the **topological charge**

$$Q = \int d^4x q(x)$$

related to the **winding number** of the field

- Several definitions on the lattice



Phenomenology of the topological charge

Topological charge is present in QCD lagrangian

$$\mathcal{L} = \frac{1}{4}F_{\mu\nu}F_{\mu\nu} + \theta q$$

- Strong CP problem
- $\eta - \eta'$ masses
- Instantons interaction
- Dependence of observables on Q

. . .

Phenomenology of the topological charge

Topological charge is present in QCD lagrangian



$$\mathcal{L} = \frac{1}{4} F_{\mu\nu} F_{\mu\nu} + \theta q$$

- Strong CP problem
- $\eta - \eta'$ masses
- Instantons interaction
- Dependence of observables on Q

. . .

Axion phenomenology - Susceptibility of the topological charge

From arXiv:1512.06746, C.Bonati, M.D'Elia, M.Mariti, G.Martinelli, M.Mesiti, F.Negro, F.S, G.Villadoro



Topological charge on the lattice

Many possibilities

Overlap operator through index theorem

Wilson Flow and field theoretical definition $Q = F_{\mu\nu}\tilde{F}_{\mu\nu}$

Cooling discrete Wilson Flow [as shown by C.Bonati, M.D'Elia PRD89 (2014)]

Stout Wilson Flow with a simpler integration scheme

Topological charge on the lattice

Many possibilities

Overlap operator through index theorem

Wilson Flow and field theoretical definition $Q = F_{\mu\nu}\tilde{F}_{\mu\nu}$


Cooling discrete Wilson Flow [as shown by C.Bonati, M.D'Elia PRD89 (2014)]

Stout Wilson Flow with a simpler integration scheme

One bottom line

They are all
equivalent
in the
continuum limit

[K.Chichy et al, Lattice'14]



- Different definitions are subject to different cut-off effects
- They can come handy for different reasons...

The history of Topological Charge Slowing Down

1996 - the prophecy: "Hybrid Monte Carlo and topological modes of full QCD"

B. Alles, G. Boyd, M. D'Elia, A. Di Giacomo and E. Vicari, Phys.Lett.B389 (1996)

"The hybrid Monte Carlo algorithm has **serious problems decorrelating the global topological charge** [...]. This represents a **warning** which must be seriously considered."

The history of Topological Charge Slowing Down

1996 - the prophecy: "Hybrid Monte Carlo and topological modes of full QCD"

B. Alles, G. Boyd, M. D'Elia, A. Di Giacomo and E. Vicari, Phys.Lett.B389 (1996)

"The hybrid Monte Carlo algorithm has **serious problems decorrelating the global topological charge** [...]. This represents a **warning** which must be seriously considered."

The message did not get across...

The history of Topological Charge Slowing Down


1996 - the prophecy: “*Hybrid Monte Carlo and topological modes of full QCD*”

B. Alles, G. Boyd, M. D’Elia, A. Di Giacomo and E. Vicari, Phys.Lett.B389 (1996)

“The hybrid Monte Carlo algorithm has **serious problems decorrelating the global topological charge** [...]. This represents a **warning** which must be seriously considered.”

The message did not get across...

2002: “ *θ dependence of $SU(N)$ gauge theories*”, L.Del Debbio, H.Panagopoulos, E.Vicari



- Non-gaussianity: $\tau_Q \neq a^{-2}$ is shown for topological charge, at variance with “ordinary” observables
- Empirical behavior: $\tau_Q \sim \exp[-c/a]$ is proposed
- Emergence of “*sizable free-energy barriers separating different regions of the configuration space*” is suggested

The history of Topological Charge Slowing Down


1996 - the prophecy: "*Hybrid Monte Carlo and topological modes of full QCD*"

B. Alles, G. Boyd, M. D'Elia, A. Di Giacomo and E. Vicari, Phys.Lett.B389 (1996)

"The hybrid Monte Carlo algorithm has **serious problems decorrelating the global topological charge** [...]. This represents a **warning** which must be seriously considered."

The message did not get across...

2002: " *θ dependence of $SU(N)$ gauge theories*", L.Del Debbio, H.Panagopoulos, E.Vicari



- Non-gaussianity: $\tau_Q \neq a^{-2}$ is shown for topological charge, at variance with "ordinary" observables
- Empirical behavior: $\tau_Q \sim \exp[-c/a]$ is proposed
- Emergence of "*sizable free-energy barriers separating different regions of the configuration space*" is suggested

2010 - the revelation: "*Properties & uses of the Wilson flow in LQCD*", M.Lüscher





At the Lattice Conference in Sardinia, The Truth is proclaimed:

- Wilson Flow should be used to define the topological charge
- In this way: "*the emergence of the topological (instanton) sectors in the continuum limit becomes transparent*"



Topological charge slowing down - two examples

Pisa Collaboration: Staggered simulations for Axion Phenomenology





Topological charge slowing down - two examples

Pisa Collaboration: Staggered simulations for Axion Phenomenology



RBC/UKQCD: Domain Wall simulations for Charm physics



“Slow Down yourself, please! Why does the system tunnel in the first place?”

A PUZZLING QUESTION

¿

How can the system
tunnel through **disconnected**
subsets of configurations?

?

“Slow Down yourself, please! Why does the system tunnel in the first place?”

A PUZZLING QUESTION

¿

How can the system
tunnel through **disconnected**
subsets of configurations?

?

Topological sectors are ill defined at finite lattice spacing

- Assignment of a configuration not unique, depends on the method
- The assignment gets clearer and clearer as we proceed to the continuum limit
- Some configurations are **particularly difficult** to assign

“Slow Down yourself, please! Why does the system tunnel in the first place?”

A PUZZLING QUESTION

?

How can the system
tunnel through **disconnected**
subsets of configurations?

?

Topological sectors are ill defined at finite lattice spacing

- Assignment of a configuration not unique, depends on the method
- The assignment gets clearer and clearer as we proceed to the continuum limit
- Some configurations are **particularly difficult** to assign

The tunneling mechanism

The system can tunnel by passing through **topologically ill defined configurations**

“Slow Down yourself, please! Why does the system tunnel in the first place?”

A PUZZLING QUESTION

?

How can the system
tunnel through **disconnected**
subsets of configurations?

?

Topological sectors are ill defined at finite lattice spacing

- Assignment of a configuration not unique, depends on the method
- The assignment gets clearer and clearer as we proceed to the continuum limit
- Some configurations are **particularly difficult** to assign

The tunneling mechanism

The system can tunnel by passing through **topologically ill defined configurations**

In the continuum limit the “breaches” are closed


- Topologically ill-defined configurations have zero measure in the continuum limit
- The tunneling gets more and more suppressed as we proceed towards the continuum

Do we have to bother?

Can't we just ignore the problem?

Do we have to bother?

Can't we just ignore the problem?




NO!

[see e.g. M.D'Elia, F.Negro, arXiv:1306.2919]

- At finite volume, Observables depends on Q
- Bad sampling of Q means to bias observables

Do we have to bother?

Can't we just ignore the problem?



NO!

[see e.g. M.D'Elia, F.Negro, arXiv:1306.2919]

- At finite volume, Observables depends on Q
- Bad sampling of Q means to bias observables

Can't we get rid of topological sectors?

2011: "Lattice QCD without topology barriers", M.Lüscher, S.Schaefer

Open the boundary on one side of the lattice

- ✓ Topological objects free to flow in/out from the system
- ✗ Loose translation invariance
- ✗ Volume effects are expected to be minimized with periodic boundaries

Simulate at strictly fixed topology

[JLQCD, PRD74 (2006)]

- ✓ In the infinite volume limit, all sectors are equivalent
- ✗ Nontrivial effects at finite volume, e.g:

$$\lim_{m \rightarrow 0} \langle q\bar{q} \rangle_{Q=0} = 0 \quad \text{at fixed } V$$

[H. Leutwyler and A. Smilga, PRD46(12) 1996]

Simulate at strictly fixed topology

[JLQCD, PRD74 (2006)]

- ✓ In the infinite volume limit, all sectors are equivalent
- ✗ Nontrivial effects at finite volume, e.g:

$$\lim_{m \rightarrow 0} \langle q \bar{q} \rangle_{Q=0} = 0 \quad \text{at fixed } V$$

[H. Leutwyler and A. Smilga, PRD46(12) 1996]

Encourage tunneling on the point x^* where the $|q(x)|$ is the largest

$$\Delta S = -\alpha \exp [-q^2(x^*)]$$

[P.de Forcrand et al., Nucl.Phys.Proc.Suppl. 63 (1998)]

- ✓ Change the action by a $1/V$ term (no need to reweight the simulation)
- ✗ Not addressing large scale topological structures

Simulate at strictly fixed topology

[JLQCD, PRD74 (2006)]

- ✓ In the infinite volume limit, all sectors are equivalent
- ✗ Nontrivial effects at finite volume, e.g:

$$\lim_{m \rightarrow 0} \langle q \bar{q} \rangle_{Q=0} = 0 \quad \text{at fixed } V$$

[H. Leutwyler and A. Smilga, PRD46(12) 1996]

Encourage tunneling on the point x^* where the $|q(x)|$ is the largest

$$\Delta S = -\alpha \exp[-q^2(x^*)]$$

[P.de Forcrand et al., Nucl.Phys.Proc.Suppl. 63 (1998)]

- ✓ Change the action by a $1/V$ term (no need to reweight the simulation)
- ✗ Not addressing large scale topological structures

Torturing the configuration by causing dislocations

“Increase the transitions rate between topological sectors by **encouraging more zero-modes**”

[G.McGlynn, R.Mawhinney, PoS lattice'13 arXiv:1311.3695]

- ? Is it working? (question for the authors in the audience)
- ? What about the scaling to the continuum?

NEW FRIENDS

CP(N-1)

MODELS

$CP(N-1)$ models in a nutshell

In the continuum - 2D space

- Commutating complex field $\vec{z} = (z_1 \dots z_N)$ of norm 1
- $U(1)$ gauge symmetry, covariant derivative: $D_\mu = \partial_\mu + iA_\mu$ with $A_\mu \in \mathcal{R}$

$$S = \beta N \int d^2x \sum_{\mu=1}^2 |D_\mu \vec{z}(x)|^2, \quad \boxed{N=21}$$

Gauge field A_μ has no kinetic term and could be integrated away, but we'd rather keep it

$CP(N-1)$ models in a nutshell

In the continuum - 2D space

- Commutating complex field $\vec{z} = (z_1 \dots z_N)$ of norm 1
- $U(1)$ gauge symmetry, covariant derivative: $D_\mu = \partial_\mu + iA_\mu$ with $A_\mu \in \mathcal{R}$

$$S = \beta N \int d^2x \sum_{\mu=1}^2 |D_\mu \vec{z}(x)|^2, \quad \boxed{N=21}$$

Gauge field A_μ has no kinetic term and could be integrated away, but we'd rather keep it

On the lattice

$$S = \beta N \sum_{n \in L^2} \sum_{\mu=1}^2 |D_\mu \vec{z}_n|^2, \quad D_\mu z_n = \Lambda_{n,\mu} z_{n+\hat{\mu}} - z_n$$

$CP(N-1)$ models in a nutshell

In the continuum - 2D space

- Commutating complex field $\vec{z} = (z_1 \dots z_N)$ of norm 1
- $U(1)$ gauge symmetry, covariant derivative: $D_\mu = \partial_\mu + iA_\mu$ with $A_\mu \in \mathcal{R}$

$$S = \beta N \int d^2x \sum_{\mu=1}^2 |D_\mu \vec{z}(x)|^2, \quad \boxed{N=21}$$

Gauge field A_μ has no kinetic term and could be integrated away, but we'd rather keep it

On the lattice

$$S = \beta N \sum_{n \in L^2} \sum_{\mu=1}^2 |D_\mu \vec{z}_n|^2, \quad D_\mu z_n = \Lambda_{n,\mu} z_{n+\hat{\mu}} - z_n$$

Like QCD...

- There is a topology Q
- There is a mass gap $M \sim 1/\xi$
- The beta-function is negative
- β sets the scale: $a \xrightarrow{\beta \rightarrow \infty} 0$

$CP(N-1)$ models in a nutshell

In the continuum - 2D space

- Commutating complex field $\vec{z} = (z_1 \dots z_N)$ of norm 1
- $U(1)$ gauge symmetry, covariant derivative: $D_\mu = \partial_\mu + iA_\mu$ with $A_\mu \in \mathcal{R}$

$$S = \beta N \int d^2x \sum_{\mu=1}^2 |D_\mu \vec{z}(x)|^2, \quad \boxed{N=21}$$

Gauge field A_μ has no kinetic term and could be integrated away, but we'd rather keep it

On the lattice

$$S = \beta N \sum_{n \in L^2} \sum_{\mu=1}^2 |D_\mu \vec{z}_n|^2, \quad D_\mu \vec{z}_n = \Lambda_{n,\mu} \vec{z}_{n+\hat{\mu}} - \vec{z}_n$$

Like QCD...

- There is a topology Q
- There is a mass gap $M \sim 1/\xi$
- The beta-function is negative
- β sets the scale: $a \xrightarrow{\beta \rightarrow \inf} 0$

But simpler!


- Simulations can be run on a laptop!
(actually: Ulisse cluster at Sissa)
- Excellent framework
to test new algorithms

MOST IMPORTANT


**it suffers
from**

**TOPOLOGICAL
FREEZING**


Topological charge evolution



Evolution on a finer lattice spacing (same scales)



Going even finer



The word "FROZEN" is rendered in a stylized, blue, icy font with a textured, crystalline appearance, set against a white background.

FROZEN

**TOPOLOGICAL
CHARGE?**




Metadynamics


Elixir

“For an *immediate relief*
of your topological ~~paralysis~~ freezing!”


Before



After the treatment




A. Laio, M. Parrinello, "Escaping free-energy minima" (2002)




Similar in spirit to Wang Landau (2001) but applied to Molecular Dynamics
Widely adopted in biochemistry (protein folding, docking, dissociation...)

**DOES
IT
WORK?**


Transition frequency vs lattice spacing - HMC



And in Metadynamics



It works at various volumes



IT WORKS!!
BUT HOW?

How does it work?

Hamiltonian dependent on **simulation time** $H(t) = H(0) + V_{bias}(t)$

How does it work?

Hamiltonian dependent on **simulation time** $H(t) = H(0) + V_{bias}(t)$

Bias potential

- V_{bias} built in terms of previous values of a **collective variable**, here taken to be Q
- Example of a possible form of the potential:

$$V_{bias}(t + dt) = V_{bias}(t) + c \cdot \exp \left[-\frac{1}{2} \left(\frac{Q - Q(t)}{\sigma} \right)^2 \right]$$

To avoid evaluating too many “exp” we actually use **triangles on a grid**

How does it work?

Hamiltonian dependent on **simulation time** $H(t) = H(0) + V_{bias}(t)$

Bias potential

- V_{bias} built in terms of previous values of a **collective variable**, here taken to be Q
- Example of a possible form of the potential:

$$V_{bias}(t + dt) = V_{bias}(t) + c \cdot \exp \left[-\frac{1}{2} \left(\frac{Q - Q(t)}{\sigma} \right)^2 \right]$$

To avoid evaluating too many “exp” we actually use **triangles on a grid**

Dynamics

- The induced force $F = -\partial_U V_{bias}$ drives the system **away** from previous values of Q
- V_{bias} **reduces** the probability of occupying previous states
- At large simulation time V_{bias} fills the free energy wells

How does it work?

Hamiltonian dependent on **simulation time** $H(t) = H(0) + V_{bias}(t)$

Bias potential

- V_{bias} built in terms of previous values of a **collective variable**, here taken to be Q
- Example of a possible form of the potential:

$$V_{bias}(t + dt) = V_{bias}(t) + c \cdot \exp \left[-\frac{1}{2} \left(\frac{Q - Q(t)}{\sigma} \right)^2 \right]$$

To avoid evaluating too many “exp” we actually use **triangles on a grid**

Dynamics


- The induced force $F = -\partial_U V_{bias}$ drives the system **away** from previous values of Q
- V_{bias} **reduces** the probability of occupying previous states
- At large simulation time V_{bias} fills the free energy wells

At convergence (long simulated time)

- V_{bias} provides a negative image of the free energy $F(Q) = -\log Z(Q)$
- The dynamics of the system is completely flat w.r.t Q

Which definition of Q ?

Geometrical: sum of the solid angle between z on all triangles



$$Q_g = \frac{1}{2\pi} \sum_{\nabla, \Delta} \arg [(\vec{z}_a, \vec{z}_b) (\vec{z}_b, \vec{z}_c) (\vec{z}_c, \vec{z}_a)]$$


This is matemagically an integer number

- ✓ **perfect** to measure the actual topological charge
- ✗ **useless** as a collective variable!

In fact $F_z = -\partial_z V_{bias}^g \propto \partial_z Q_g = 0$: the bias would induce no force on the system

Which definition of Q ?

Geometrical: sum of the solid angle between z on all triangles




$$Q_g = \frac{1}{2\pi} \sum_{\nabla, \Delta} \arg [(\vec{z}_a, \vec{z}_b) (\vec{z}_b, \vec{z}_c) (\vec{z}_c, \vec{z}_a)]$$

This is matemagically an integer number

- ✓ **perfect** to measure the actual topological charge
- ✗ **useless** as a collective variable!

In fact $F_z = -\partial_z V_{bias}^g \propto \partial_z Q_g = 0$: the bias would induce no force on the system

Gauge definition: plaquette of Λ




$$Q = \frac{1}{2\pi} \sum_{\square} \text{Im} \square = ZQ_g + \eta - \text{Not an integer number}$$

- ✗ **not ideal** to measure the actual topological charge
- ✓ **useful** as a collective variable: $F_{\Lambda} = -\partial_{\Lambda} V_{bias}^Q \propto \partial_{\Lambda} Q \neq 0$
- Field Λ must be smoothed, so that $\sqrt{\langle \eta^2 \rangle} \lesssim 1$ and $Z \sim 1$
- Analytical smoothing easily differentiable: **stout smearing**

Which definition of Q ?

Geometrical: sum of the solid angle between z on all triangles




$$Q_g = \frac{1}{2\pi} \sum_{\nabla, \Delta} \arg [(\vec{z}_a, \vec{z}_b) (\vec{z}_b, \vec{z}_c) (\vec{z}_c, \vec{z}_a)]$$

This is matemagically an integer number

- ✓ **perfect** to measure the actual topological charge
- ✗ **useless** as a collective variable!

In fact $F_z = -\partial_z V_{bias}^g \propto \partial_z Q_g = 0$: the bias would induce no force on the system


Gauge definition: plaquette of Λ





$$Q = \frac{1}{2\pi} \sum_{\square} \text{Im} \square = ZQ_g + \eta - \text{Not an integer number}$$

- ✗ **not ideal** to measure the actual topological charge
- ✓ **useful** as a collective variable: $F_{\Lambda} = -\partial_{\Lambda} V_{bias}^Q \propto \partial_{\Lambda} Q \neq 0$
- Field Λ must be smoothed, so that $\sqrt{\langle \eta^2 \rangle} \lesssim 1$ and $Z \sim 1$
- Analytical smoothing easily differentiable: **stout smearing**

What's the shape of $F(Q)$?







“What about the **sampled distribution** of Q ?”

At convergence

By construction $F(Q) = -\log Z(Q)$ which means that

$$P(Q) = \text{const}$$

in the generated sample

“What about the **sampled distribution** of Q ?”

At convergence

By construction $F(Q) = -\log Z(Q)$ which means that

$$P(Q) = \text{const}$$

in the generated sample

“So you are sampling a different distribution!!!”

$F(Q)$ can be used to **reweight** the distribution:

$$\langle O \rangle = \frac{\sum_i O_i \exp[-F(Q_i)]}{\sum_j \exp[-F(Q_j)]}$$

“What about the **sampled distribution** of Q ?”

At convergence

By construction $F(Q) = -\log Z(Q)$ which means that

$$P(Q) = \text{const}$$

in the generated sample

“So you are sampling a different distribution!!!”

$F(Q)$ can be used to **reweight** the distribution:


$$\langle O \rangle = \frac{\sum_i O_i \exp[-F(Q_i)]}{\sum_j \exp[-F(Q_j)]}$$

Reweighting costs


- By reweighting we suppress configurations with non-integer charge
- Nonetheless the configurations generated by metadynamics are uncorrelated
 - We agree with HMC where it works, but we achieve increasingly large speed-up as $a \rightarrow 0$
 - We obtain sensible results at reasonable cost, even when the HMC is completely frozen

The associated costs seems to scale well with a and V (see next plots)


$\rho(Q)$, HMC (40M painful trajectories, $\beta = 0.75$, $\xi/a \sim 5.16$, $L/a = 60$)




$\rho(Q)$, metadynamics (700k trajectories)




Reweighting



Topological susceptibility



Topological susceptibility - The cost



“The algorithm is not ergodic!”

- Left to itself, the algorithm would explore with equal probability **all topological sectors**
- We don't want to sample sectors with **too large charge** (suppressed after reweighting)
- We **constrain** the dynamic in the range $[-Q_{max}; +Q_{max}]$
- Q_{max} must be chosen to be much **larger** than $\sqrt{\langle Q^2 \rangle}$
- Q_{max} can be taken as large **as wanted**

"The algorithm is not ergodic!"

- Left to itself, the algorithm would explore with equal probability **all topological sectors**
- We don't want to sample sectors with **too large charge** (suppressed after reweighting)
- We **constrain** the dynamic in the range $[-Q_{max}; +Q_{max}]$
- Q_{max} must be chosen to be much **larger** than $\sqrt{\langle Q^2 \rangle}$
- Q_{max} can be taken as large **as wanted**

"You are violating the sacred principles of Monte Carlo methods!"

- In fact the algorithm does not build a Markov Chain of configurations $[z, \Lambda]$ at all!
- You have to think in terms of the enlarged configuration space $\{[z, \Lambda] \otimes V_{bias}\}$
- Indeed it was rigorously shown that:

The correct sampling of the configuration space is obtained

after reweighting

[Equilibrium Free Energies from Nonequilibrium Metadynamics,

G.Bussi, A.Laio, M.Parrinello, PRL96 (2006)]

No conceptual difference

It amounts to simulate with a time-dependent (imaginary) $V_{bias} = \theta_{QCD} Q^{stout}$ where

$$\theta_{QCD}(t) = i F [Q^{stout}(t)]$$

Tune the ~ 5 parameters on the basis of the $CP(N-1)$ experience

No conceptual difference

It amounts to simulate with a time-dependent (imaginary) $V_{bias} = \theta_{QCD} Q^{stout}$ where

$$\theta_{QCD}(t) = i F [Q^{stout}(t)]$$

Tune the ~ 5 parameters on the basis of the $CP(N-1)$ experience

Ingredients

- Compute a new force term $\propto \partial_U Q$
- Stout smear the configuration (several levels, $\mathcal{O}(10)$ needed)
- Remap the force iteratively $F^{non-stout} \rightarrow F^{1-stout} \rightarrow \dots F^{N-stout}$

Extension to QCD

No conceptual difference

It amounts to simulate with a time-dependent (imaginary) $V_{bias} = \theta_{QCD} Q^{stout}$ where

$$\theta_{QCD}(t) = i F [Q^{stout}(t)]$$

Tune the ~ 5 parameters on the basis of the $CP(N-1)$ experience


Ingredients

- Compute a new force term $\propto \partial_U Q$
- Stout smear the configuration (several levels, $\mathcal{O}(10)$ needed)
- Remap the force iteratively $F^{non-stout} \rightarrow F^{1-stout} \rightarrow \dots F^{N-stout}$


A first taste - In collaboration also with M.D'Elia, C.Bonati

Can we unfreeze this? \longrightarrow

- | | |
|------------------------|------------------|
| • $\beta = 4.36$ | • staggered |
| • $a = 0.0397$ fm | • $N_f = 2 + 1$ |
| • $M_\pi \sim 135$ MeV | • small volume |
| • $L/a = 40$ | • totally frozen |



It looks promising...



Squeezing the best from the algorithm

- Make use of $Q \rightarrow -Q$ symmetry
- Make use of $Q \rightarrow Q + 2k\pi$ symmetry?
- Precondition the algorithm, feeding-in the information on $F(Q)$
- Improve the convergence starting from a guess of V_{bias}
- Include other collective variables

Future improvements

Squeezing the best from the algorithm

- Make use of $Q \rightarrow -Q$ symmetry
- Make use of $Q \rightarrow Q + 2k\pi$ symmetry?
- Precondition the algorithm, feeding-in the information on $F(Q)$
- Improve the convergence starting from a guess of V_{bias}
- Include other collective variables

Extending to QCD

- No conceptual problems, just a bit of pain to implement
- Preliminary test shows encouraging results
- Needs more stout: 30-40% overhead (less important towards the continuum limit)

Future improvements

Squeezing the best from the algorithm

- Make use of $Q \rightarrow -Q$ symmetry
- Make use of $Q \rightarrow Q + 2k\pi$ symmetry?
- Precondition the algorithm, feeding-in the information on $F(Q)$
- Improve the convergence starting from a guess of V_{bias}
- Include other collective variables

Extending to QCD

- No conceptual problems, just a bit of pain to implement
- Preliminary test shows encouraging results
- Needs more stout: 30-40% overhead (less important towards the continuum limit)

More than topology?

- Can it be used to study **Gribov copies** problem in Gauge Fixing?
- Can it help computing **Spectral Density**?
- Can it be used to study **Finite Density**!?

Conclusions

Topology

- Different definitions of the Topological charge can be **useful for different reasons**
- Dependency on the topological sector is **non trivial**
- Simulations get frozen close to the continuum limit (**a long history**)

Metadynamics

Coupling **the past** history to reduce the occupancy of already explored states

- Bias potential inducing a force driving “**away from the past**”
- Topological charge gets **unfrozen**
- Distribution of Q at Long Simulation Time is **flat**: $P(Q) = 1$
- Reweighting restores the proper distribution
- Several parameters to tune...

The future

- Use all the available **symmetries**
- Further test **QCD** simulations
- Apply to **other problems**

...THANKS...



**...FOR YOUR
ATTENTION!!!**